


UNIVERSIDADE FEDERAL DO PARANÁ

GUSTAVO HENRIQUE QUEIROZ SCHUNEMANN MANFRIN DE OLIVEIRA

EFEITOS DA GALANTAMINA NO COMPROMETIMENTO DA MEMÓRIA E  
AUMENTO DE BETA-AMILÓIDE E ALFA-SINUCLÉINA NO MODELO ANIMAL  
DE DOENÇA DE ALZHEIMER INDUZIDO POR ESTREPTOZOTOCINA.

CURITIBA

2020



GUSTAVO HENRIQUE QUEIROZ SCHUNEMANN MANFRIN DE OLIVEIRA

EFEITOS DA GALANTAMINA NO COMPROMETIMENTO DA MEMÓRIA E  
AUMENTO DE BETA-AMILÓIDE E ALFA-SINUCLEÍNA NO MODELO ANIMAL  
DE DOENÇA DE ALZHEIMER INDUZIDO POR ESTREPTOZOTOCINA.

Dissertação apresentada como requisito  
parcial à obtenção do grau de Mestre em  
Farmacologia, no Curso de Pós-  
Graduação em Farmacologia, Setor de  
Ciências Biológicas, Universidade  
Federal do Paraná.

Orientadora: Profa. Dra. Maria Aparecida  
Barbato Frazao Vital.

CURITIBA

2020

Universidade Federal do Paraná  
Sistema de Bibliotecas  
(Giana Mara Seniski Silva – CRB/9 1406)

Oliveria, Gustavo Henrique Queiroz Schunemann Manfrin de  
Efeitos de galantamina no comprometimento da memória e aumento de  
beta-amilóide e alfa-sinucleína no modelo animal de doença de Alzheimer  
induzido por estreptozotocina. / Gustavo Henrique Queiroz Schunemann  
Manfrin de Oliveira. – Curitiba, 2020.  
20 p.: il.

Orientadora: Maria Aparecida Barbato Frazão Vital.

Dissertação (mestrado) - Universidade Federal do Paraná, Setor de  
Ciências Biológicas. Programa de Pós-Graduação em Farmacologia.

1. Alzheimer, Doença de. 2. Galantamina. 3. Estreptozocina. 4. alfa-  
Sinucleína. 5. Peptídeos beta-Amiloides. 6. Memória. I. Título. II. Vital,  
Maria Aparecida Barbato Frazão, 1964-. III. IV. Universidade Federal do  
Paraná. Setor de Ciências Biológicas. Programa de Pós-Graduação em  
Farmacologia.

CDD (22. ed.) 616.831



MINISTÉRIO DA EDUCAÇÃO  
SETOR DE CIÊNCIAS BIOLÓGICAS  
UNIVERSIDADE FEDERAL DO PARANÁ  
PRÓ-REITORIA DE PESQUISA E PÓS-GRADUAÇÃO  
PROGRAMA DE PÓS-GRADUAÇÃO FARMACOLOGIA -  
40001016038P0

## TERMO DE APROVAÇÃO

Os membros da Banca Examinadora designada pelo Colegiado do Programa de Pós-Graduação em FARMACOLOGIA da Universidade Federal do Paraná foram convocados para realizar a arguição da Dissertação de Mestrado de **GUSTAVO HENRIQUE QUEIROZ SCHUNEMANN MANFRIN DE OLIVEIRA** intitulada: **EFEITOS DA GALANTAMINA NO COMPROMETIMENTO DA MEMÓRIA E AUMENTO DE ETA-AMILÓIDE E ALFA-SINUCLEÍNA NO MODELO ANIMAL DE DOENÇA DE ALZHEIMER INDUZIDO POR ESTREPTOZOTOCINA**, sob orientação da Profa. Dra. MARIA APARECIDA BARBATO FRAZÃO VITAL, que após terem inquirido o aluno e realizada a avaliação do trabalho, são de parecer pela sua APROVAÇÃO no rito de defesa. A outorga do título de mestre está sujeita à homologação pelo colegiado, ao atendimento de todas as indicações e correções solicitadas pela banca e ao pleno atendimento das demandas regimentais do Programa de Pós-Graduação.

CURITIBA, 15 de Outubro de 2020.

Assinatura Eletrônica

20/10/2020 14:54:03.0

MARIA APARECIDA BARBATO FRAZÃO VITAL

Presidente da Banca Examinadora (UNIVERSIDADE FEDERAL DO PARANÁ)

Assinatura Eletrônica

20/10/2020 09:01:18.0

ROBERTO ANDREATINI

Avaliador Interno (UNIVERSIDADE FEDERAL DO PARANÁ)

Assinatura Eletrônica

20/10/2020 11:31:29.0

DEBORA DALLA VECCHIA

Avaliador Externo (FACULDADE PARANAENSE)

---

Centro Politécnico - CURITIBA - Paraná - Brasil

CEP 81531990 - Tel: (0xx41)3361-1003 - E-mail: pgfarmacologia@ufpr.br

Documento assinado eletronicamente de acordo com o disposto na legislação federal Decreto 8539 de 08 de outubro de 2015.

Gerado e autenticado pelo SIGA-UFPR, com a seguinte identificação única: 56508

Para autenticar este documento/assinatura, acesse <https://www.prppg.ufpr.br/siga/visitante/autenticacaoassinaturas.jsp> e insira o código 56508

## **NOTA EXPLICATIVA**

Esta dissertação é apresentada em formato alternativo – artigo para publicação – de acordo com as normas do Programa de Pós-Graduação em Farmacologia da Universidade Federal do Paraná.

## RESUMO

O modelo de doença de Alzheimer (DA) induzida por Estreptozotocina (STZ) gera um estado de resistência cerebral à insulina, capaz de causar prejuízo cognitivo nos animais que se assemelha aquele observado na DA, devido ao estresse oxidativo mediado por processo inflamatório. Os níveis de citocinas inflamatórias podem ser reduzidos pelo tratamento com Galantamina, uma vez que esse fármaco pode interferir no metabolismo glicídico; assim, seu uso é indicado para amenizar os sintomas da DA uma vez que a deficiência no metabolismo da glicose pode participar da patogênese da DA. Neste contexto, foram investigados os efeitos da Galantamina na memória e frequência de locomoção de ratos Wistar tratados com STZ. Para isso, mimetizamos a DA em ratos Wistar machos por meio de injeções intracerebroventriculares de STZ (1mg / kg). Em seguida, testes comportamentais foram realizados em 32 ratos: 16 injetados com STZ e 16 injetados com veículo. Posteriormente, metade dos ratos em cada um dos grupos foi tratado com Galantamina (5 mg / kg). A outra metade recebeu veículo. Além disso, realizamos teste tipo ELISA para avaliar o acúmulo de Alfa Sinucleína ( $\alpha$ -Syn) e Beta Amilóide ( $A\beta$ ) nos cérebros dos ratos. Os resultados de alguns testes comportamentais indicaram que o grupo STZ + Veículo possuía prejuízos cognitivos, quando comparado ao grupo Sham + Veículo. Não houveram diferenças estatisticamente significantes na memória dos ratos tratados com Galantamina, quando comparados aos tratados apenas com veículo. O ELISA mostrou acúmulo de  $A\beta$  e  $\alpha$ -Syn em grupos de ratos lesados pela STZ. Este último resultado sugere que o comprometimento da locomoção encontrado nos animais pode ser devido ao acúmulo de  $\alpha$ -Syn e não apenas pela mimetização de DA causada pela STZ.

Palavras-chave: Doença de Alzheimer. Galantamina. Estreptozotocina. Alfa Sinucleína. Beta Amilóide.



## ABSTRACT

The Streptozotocin (STZ) induced animal model of Alzheimer's Disease (AD) generates a state of cerebral insulin resistance, capable of mimicking AD by causing oxidative stress mediated by inflammatory process. The levels of inflammatory cytokines can be reduced by Galantamine treatment; thus, its use is indicated to ease AD symptoms, since it is believed that deficiency in glucose metabolism may participate in the pathogenesis of AD. In this context, we aim to investigate the effects of Galantamine regarding memory and locomotion frequency in Wistar rats treated with STZ. To do so, we mimic the AD in male Wistar rats through intracerebroventricular injections of STZ (1mg/kg). Then, behavioral testes were carried for 32 rats: 16 injected with STZ and 16 injected with vehicle. Subsequently, we treated half of the rats in each of the aforementioned groups with Galantamine (5 mg/kg). The other half of the rats of each group were given vehicle. In addition, we performed ELISA to assess Beta Amyloid ( $A\beta$ ) and Alpha Synuclein ( $\alpha$ -Syn) accumulation in the rats' brains. Except for the Novel Object Recognition Test, all other behavioral tests results indicated that the STZ + Vehicle group was impaired when compared to the Sham + Vehicle group. In addition, our results showed no statistically significant differences in the memory of the rats treated with Galantamine, when compared to those only treated with vehicle, regarding behavioral tests. The ELISA showed  $A\beta$  and  $\alpha$ -Syn accumulation in some groups of rats impaired with STZ. This last result suggests that the locomotion impairment found in the animals could be due to the  $\alpha$ -Syn accumulation and not only because of the AD mimicking caused by the STZ. Finally, the animals impaired with STZ did not presented improvements when treated with Galantamine, nonetheless, the overall performance for the STZ + Galantamine group was higher than the STZ + Vehicle one on the NORT. Hence, we believe that more studies on this subject should be carried out to elucidate some challenges yet unresolved.

Keywords: Alzheimer's Disease, Galantamine, Streptozotocin, Alpha Synuclein, Beta Amyloid.

## SUMÁRIO

<b>1 INTRODUÇÃO .....</b>	<b>16</b>
<b>2 OBJETIVOS .....</b>	<b>19</b>
2.1 OBJETIVO GERAL .....	19
2.2 OBJETIVOS ESPECÍFICOS .....	19
<b>3 ARTIGO CIENTÍFICO .....</b>	<b>16</b>
<b>4 CONSIDERAÇÕES FINAIS .....</b>	<b>16</b>
<b>5 CONCLUSÃO .....</b>	<b>17</b>
<b>REFERÊNCIAS.....</b>	<b>18</b>



## 1 INTRODUÇÃO

Demência é a principal causa de distúrbios progressivos em funções cognitivas como linguagem, memória, pensamento racional e habilidades sociais, além disso, ela é capaz de promover alterações emocionais e de personalidade (Cipriani *et al.*, 2015). Um dos tipos mais comuns de demência é a doença de Alzheimer (DA). Atualmente, estima-se que 35,6 milhões de pessoas vivam com ela e acredita-se que essa estimativa possa dobrar a cada 20 anos, chegando a 65,7 milhões em 2030 (Dos Santos *et al.*, 2018).

Por ser uma doença neurodegenerativa progressiva de caráter irreversível, a DA afeta a cognição e interfere na memória, o que leva o indivíduo à demência total em estágios avançados. No entanto, a doença também pode interferir nas capacidades físicas de pessoas afetadas (memória, cognição, fala e funções motoras), o que a torna a principal causa de dependência funcional, institucionalização e mortalidade na população idosa (Dos Santos *et al.*, 2018; Silvestrelli *et al.*, 2006).

A neuropatologia da DA segue um padrão semelhante a outras demências, existindo um certo arquétipo seguido pelas células neurais que são acometidas: lesão neuronal, falha sináptica e finalmente morte celular (Kamat *et al.*, 2016; Lowe *et al.*, 2018).

Esses achados levam às duas hipóteses principais de alterações cerebrais causadas pela DA: a hipótese da cascata amilóide e a hipótese colinérgica. Ambas as hipóteses sugerem que o desfecho final da DA é a perda funcional com a consequente morte neuronal, pelo acúmulo de proteína Beta Amilóide (A $\beta$ ) e emaranhados TAU (Barage e Sonawane, 2015; Pákási e Kálmán, 2008; Sereniki e Vital, 2008).

Outra proteína potencialmente nociva é a  $\alpha$ -sinucleína ( $\alpha$ -Syn). Seu acúmulo pode levar a um processo de envelhecimento errôneo, através de corpos de Lewy, culminando em doenças neurodegenerativas como a doença de Parkinson (DP) e a doença do corpo de Lewy (Spillantini e Dickson, 2003). Embora a  $\alpha$ -Syn seja comumente considerada um biomarcador de DP (Chang *et al.*, 2020), alguns estudos

demonstram que até 60% dos pacientes com DA apresentam sinais de distúrbios causados por corpos de Lewy (Arai *et al.*, 2001; Hamilton, 2000; James *et al.*, 2012; Schneider *et al.*, 2009). Em um estudo de coorte com 225 pacientes com DA, foi demonstrado que a concentração de  $\alpha$ -Syn no líquido cefalorraquidiano estava significativamente aumentada (Majbour *et al.*, 2017). Outro estudo sugere que os distúrbios da DA tenham ligação com lesões simultâneas mediados por  $\alpha$ -Syn (Mikolaenko *et al.*, 2005). Apesar de serem diferentes os possíveis desfechos clínicos oriundo dos acúmulo de A $\beta$  e  $\alpha$ -Syn, ambas proteínas compartilham mecanismos que levam ao estresse oxidativo (Angelova e Abramov, 2017).

Estudos sugerem que a diminuição do metabolismo neural da glicose desencadeia a cascata neurodegenerativa da DA (De La Monte e Wands, 2008; Grieb, 2016; Steen *et al.*, 2005). A partir disso, a Estreptozotocina (STZ), uma droga capaz de induzir o estado cerebral resistente à insulina (IRBS) (Barilar *et al.*, 2020), pode ser usada para mimetizar algumas características clínicas da DA, quando administrada por via intracerebroventricular (ICV-STZ).

Apesar da literatura evidenciar conexões entre a proteína A $\beta$  e a DA, ainda é necessário um melhor entendimento da relação entre a DA e a  $\alpha$ -Syn. Até onde os autores desse trabalho tem conhecimento, não foram encontrados na literatura estudos sobre a relação entre a STZ, A $\beta$  e  $\alpha$ -Syn. Dessa forma, decidimos por investigar este assunto de maneira mais aprofundada neste trabalho.

Um crescente acúmulo de evidências sugere efeito neuroprotetor da Galantamina (Bloniecki *et al.*, 2017; Geerts, 2005; Geerts *et al.*, 2005; Wang *et al.*, 2019). A Galantamina possui propriedades antioxidantes, antiapoptóticas e antiinflamatórias, de tal maneira que pode modular o metabolismo da glicose (Ali, 2015; El-Abhar *et al.*, 2015).

A dose comumente usada em ratos varia de 1 mg / kg a 5 mg / kg (Furukawa *et al.*, 2014; Geerts, 2005; Geerts *et al.*, 2005; Satapathy *et al.*, 2011). Nesse cenário, um estudo demonstrou que ratos recém-nascidos pré-tratados com Galantamina tiveram redução nos danos cerebrais induzidos por hipóxia-isquemia. Neste estudo, a dose de 5 mg / kg de Galantamina mostrou acentuada diminuição de danos cerebrais quando comparada ao grupo controle e aos grupos pré-tratados com 1 e 2,5 mg / kg de Galantamina (Furukawa *et al.*, 2014).

Embora muito já se tenha sido elucidado acerca desse assunto, mais estudos sobre os efeitos farmacodinâmicos da Galantamina são necessários para apoiar a evidência de que ela pode ser usada efetivamente como um tratamento para os sintomas da DA (El-Abhar *et al.*, 2015). Portanto, o objetivo deste estudo foi analisar os efeitos da Galantamina em ratos por meio de um modelo animal da DA induzida por STZ.

Para avaliar os efeitos da Galantamina, primeiramente propomos mimetizar a AD por meio de ICV-STZ em ratos. Então, realizando os testes comportamentais Labirinto Y, Teste de Campo Aberto e Teste de Reconhecimento de Novo Objeto em grupos de animais, é possível medir os efeitos da STZ e do tratamento com Galantamina nos animais. Para analisar as implicações desses fármacos a nível bioquímico, o hipocampo e o córtex pré-frontal dos animais foram analisados pelo método *enzyme-linked immunosorbent assay* (ELISA), a fim de se quantificar as proteínas A $\beta$  e  $\alpha$ -Syn.

## 2 OBJETIVOS

### 2.1 OBJETIVO GERAL

Investigar os efeitos da Galantamina em um modelo animal de DA induzido por STZ.

### 2.2 OBJETIVOS ESPECÍFICOS

Mimetizar a DA através de ICV-STZ em uma amostra aleatória de ratos

Realizar os testes comportamentais Labirinto em Y, Teste de Campo Aberto e Reconhecimento de Novos Objetos; para medir os efeitos da STZ e implicações do tratamento com a Galantamina.

Analisar o Hipocampo e o cortex pré-frontal através do método ELISA, para quantificar  $A\beta$  e  $\alpha$ -Syn.

Relacionar os efeitos do tratamento farmacológico com os testes comportamentais e análises bioquímicas.

### 3 ARTIGO CIENTÍFICO

#### **Effects of Galantamine in Memory Impairment and Increase of $\beta$ -Amyloid and Alpha-synuclein in Streptozotocin Induced an Animal Model of Alzheimer's Disease.**

Gustavo S. Oliveira <sup>a\*</sup>, Leonardo C Souza <sup>a</sup>, Evellyn M. Azevedo <sup>b</sup>, Guilherme Q. Oliveira <sup>c</sup>, Maria A. B. F. Vital <sup>a\*</sup>

<sup>a</sup> Department of Pharmacology, Federal University of Paraná, PR, Brazil.

<sup>b</sup> Department Physiology, Federal University of Paraná, PR, Brazil.

<sup>c</sup> Engenharia Mecânica, Escola Politécnica, Universidade Positivo, PR, Brazil.

\*Corresponding authors:

E-mail address: [gschunemann@gmail.com](mailto:gschunemann@gmail.com) (Gustavo S. Oliveira); [vital@ufpr.br](mailto:vital@ufpr.br) (M.A.B.F. Vital).

## ABSTRACT

The Streptozotocin (STZ) induced animal model of Alzheimer's Disease (AD) generates a state of cerebral insulin resistance, capable of mimicking AD by causing oxidative stress mediated by inflammatory process. The levels of inflammatory cytokines can be reduced by Galantamine treatment; thus, its use is indicated to ease AD symptoms, since it is believed that deficiency in glucose metabolism may participate in the pathogenesis of AD. In this context, we aim to investigate the effects of Galantamine regarding memory and locomotion frequency in Wistar rats treated with STZ. To do so, we mimic the AD in male Wistar rats through intracerebroventricular injections of STZ (1mg/kg). Then, behavioral testes were carried for 32 rats: 16 injected with STZ and 16 injected with vehicle. Subsequently, we treated half of the rats in each of the aforementioned groups with Galantamine (5 mg/kg). The other half of the rats of each group were given vehicle. In addition, we performed ELISA to assess Beta Amyloid ( $A\beta$ ) and Alpha Synuclein ( $\alpha$ -Syn) accumulation in the rats' brains. Except for the Novel Object Recognition Test, all other behavioral tests results indicated that the STZ + Vehicle group was impaired when compared to the Sham + Vehicle group. In addition, our results showed no statistically significant differences in the memory of the rats treated with Galantamine, when compared to those only treated with vehicle, regarding behavioral tests. The ELISA showed  $A\beta$  and  $\alpha$ -Syn accumulation in some groups of rats impaired with STZ. This last result suggests that the locomotion impairment found in the animals could be due to the  $\alpha$ -Syn accumulation and not only because of the AD mimicking caused by the STZ. Finally, the animals impaired with STZ did not presented improvements when treated with Galantamine, nonetheless, the overall performance for the STZ + Galantamine group was higher than the STZ + Vehicle one on the NORT. Hence, we believe that more studies on this subject should be carried out to elucidate some challenges yet unresolved.

**Keywords:** Alzheimer's Disease, Galantamine, Streptozotocin, Alpha Synuclein, Beta Amyloid.

## 1. Introduction

Dementia is the main cause of progressive disorders in cognitive functions, such as language, memory, rational thinking and social skills, as well as changes in emotion and personality [1]. One of the most common types of dementia is the well-known Alzheimer's disease (AD). Currently, it is estimated that 35.6 million people live with AD and it is believed that this estimate may double every 20 years, reaching 65.7 million in 2030 [2]. As an irreversible and progressive disease AD affects cognition, mainly affecting memory, which leads the individual to total dementia in advanced stages. Nevertheless, the disease can also involve the motor capabilities of those affected by this pathology which makes it the main cause of functional dependence, institutionalization and mortality among the elderly population [2, 3].

The neuropathology of AD follows a pattern similar to other dementias. There a pattern to be followed by neural cells in which the disease develops: neuronal injury, synaptic failure and finally cell death [4, 5]. These findings lead to the two main hypotheses of brain changes caused by AD: the amyloid cascade hypothesis and the cholinergic hypothesis. Both hypotheses suggest that the final outcome of the impairment caused by the AD cascade is the functional loss with the consequent neuronal death, by the accumulation of Beta amyloid protein ( $A\beta$ ) and TAU tangles [6-8].

In addition, another potential harmful protein is  $\alpha$ -synuclein ( $\alpha$ -Syn). Its lesions may lead to an unhealthy aging process, through Lewy body, ending up in neurodegenerative disorders such as Parkinson's disease (PD) and Lewy Body Disease [9]. Although  $\alpha$ -Syn is commonly considered as a PD biomarker [10], some studies demonstrate that up to 60% of patients with AD have signs of Lewy body disorders [11-14]. In a cohort study with 225 AD patients, it was shown that the  $\alpha$ -Syn concentration on the cerebrospinal fluid was significantly increased [15]. Other study suggests that the disturbances of the AD has linkage with simultaneous  $\alpha$ -Syn lesions [16]. Moreover, despite the key targets of  $A\beta$  and  $\alpha$ -Syn are dissimilar, either share mechanisms that leads to oxidative stress [17].

Studies suggest that the decrease in neural glucose metabolism triggers the neurodegenerative cascade of AD [18-20]. From this, Streptozotocin (STZ), a drug



able to induce Insulin Resistant Brain State (IRBS) [21], can be used to mimic some AD clinical features when administered intracerebroventricularly (ICV-STZ). Through low dose injections of STZ into the lateral ventricles of animal's brains, this drug induces oxidative stress [22], neuroinflammation [23], damage to the cerebral cholinergic system [24], formation of positive aggregates for protein  $A\beta$  [25] and impairments in cerebral energy metabolism [26]; resulting in cognitive deficits such as impaired memory and spatial orientation [27].

Despite these previously mentioned connections between  $\alpha$ -Syn and  $A\beta$  regarding the AD, a better understanding of the relationship between AD and  $\alpha$ -Syn is needed. Hence, and since to the best of the authors knowledge no study concerning the relation of STZ,  $A\beta$  and  $\alpha$ -Syn were found in the literature, we decided to investigate this matter a little further in this paper.

Several studies have accumulated evidence that suggests the neuroprotective effect of Galantamine [28-30]. Galantamine has anti-oxidant, anti-apoptotic and anti-inflammatory properties, that way, through its properties it could modulate glucose metabolism [31-33]. The dose commonly used by researchers varies from 1 mg/kg to 5 mg/kg [29, 34-36]. In this scenario, a study found that newborn rats pre-treated with Galantamine have reduced brain damage induced by hypoxia-ischemia. In this study the 5 mg/kg dose of Galantamine showed marked reduction of brain damage when compared to the control group and the 1 and 2.5 mg/kg Galantamine pre-treated groups [34].

However, further studies regarding the pharmacodynamic effects of the Galantamine are required to support the evidence that it can be used effectively as a treatment to the AD symptoms [31]. Therefore, this paper aims to analyze the effects of Galantamine on rats with STZ induced animal model of AD.

To evaluate the Galantamine's effects, first we propose to mimic the AD through ICZ-STZ in the rats. Then, by performing the Y Maze, Open Field Test (OFT) and Novel Object Recognition Test (NORT) on an animal sample, it is possible to measure the STZ and the Galantamine treatment impacts on the animal's performance. Here, the hippocampus and pre-frontal cortex of the animals were also analyzed by enzyme-linked immunosorbent assay (ELISA), to quantify the  $A\beta$  and  $\alpha$ -Syn.

## **2. Methods**

### **2.1. Animals**

Male Wistar rats, approximately 3 months old, were used at the beginning of the experiments, from the vivarium bioterium of the Biological Sciences Sector of the Federal University of Paraná. The rats were kept in groups of a maximum of 5 per box, in a room with controlled humidity and temperature ( $22 \pm 2^\circ \text{C}$ ), a 12-hour light-dark cycle, with the light phase starting at 7:00 am. Water and food were provided at will during all experiments.


All procedures were conducted according to the ethics and research protocols of the Federal University of Paraná and this research was approved by the university's ethics committee, with the number 1226.

### **2.2. Drugs**

- Streptozotocin 1 mg/kg (STZ) was purchased from Santa Cruz Biotechnology Inc., Santa Cruz, USA.
- Galantamine 5 mg/kg was purchased from Prati, Donaduzzi & Cia Ltda., Toledo, Brazil.

### **2.3. Experimental design**

Thirty-two rats were randomly divided into groups sham ( $n = 16$ ) and STZ ( $n = 16$ ). STZ group received bilateral injection of STZ, at a dose of 1 mg/kg, dissolved in 0.9% sterile saline solution in the lateral ventricles. The sham group received only sterile saline. Daily administrations of Galantamine were performed to assess the effects of this treatment in STZ injured animals, on cognitive functions, spatial memory, locomotion frequency and short-term recognition. For such treatment, the sham and STZ groups were randomly subdivided into 4 groups of 8 animals each: Sham + Vehicle; Sham + Galantamine; STZ + Vehicle and STZ + Galantamine (Figure 1).




**Figure 1** - Groups, procedures and number of animals per group. ICV: Intracerebroventricular; ICV-STZ: Intracerebroventricular Streptozotocin.

Galantamine was administered to the animals for 30 days intraperitoneally. It started 1h before stereotaxic surgery, using 0.5% carboxymethylcellulose in distilled water with 1% Tween 80 as a vehicle. Galantamine was able to produce results in Rats in doses of 4 mg/kg or 5 mg/kg [36]. In this scenario, the dose of Galantamine used in this paper was 5 mg/kg. Behavioral tests were performed on the animals:

- On the 21st day after surgery, the OFT was performed.
- On the 28th day the rats went through the habituation of the box of objects and then on the 29th day they performed the object location test.
- On the 30th day, the Y Maze test was fulfilled.

At the end of the behavioral tests, on the 30th day of the experiment, the animals were euthanized and the brains dissected for measurement of  $\beta$ -amyloid protein and alpha synuclein in the hippocampus and prefrontal cortex.



**Figure 2** – Experimental design. Scheme of procedures and tests performed. ICV: Intracerebroventricular; ICV-STZ: Intracerebroventricular Streptozotocin OFT: Open Field Test; NORT: Novel Object Recognition Test.

## **2.4. Stereotaxic surgery**

Before stereotaxic surgery, the animals were anesthetized with Equitesin 0.3mg/kg (distilled water, chloral hydrate, magnesium sulfate, 90% alcohol, thiopental and propylene glycol). Thereafter, they received atropine sulfate (0.4 mg / kg, i.p.). After this procedure, the animals were placed in a stereotaxic (David Kopf, model 957L). Through a scalpel, access to the skull was established and for the location of the ventricles, the following coordinates were adjusted: anteroposterior -0.8 mm, lateral 1.5 mm and dorsoventral 4.0 mm relative to the bregma and ventral from the dura with the bar set to 0 mm [37]. After craniotomy, STZ was infused with a 30-gauge needle connected to a polyethylene tube connected to a 10 $\mu$ l micro-syringe (Hamilton, USA). The composition of this equipment was linked to an infusion pump (Havard Apparatus, USA). In the STZ groups, 1mg / kg of STZ (5  $\mu$ l per injection site) was infused into the lateral ventricles. In the sham groups, the same methodology was used, however, the rats were treated with sterile saline solution, instead of STZ.

At the end of the surgeries, the rats were transferred to individual boxes in a heated room, where they were able to recover from the procedure and only then could be transported to vivarium.

## **2.5. Behavioral Evaluation**

### **2.5.1 Open Field Test**

On the 21st day after the surgery, the OFT was performed to analyze spontaneous locomotor activity. The OFT consists of a circular arena (97-cm diameter, 42-cm height) separated into three concentric circles and subdivided into 19 quadrants. Each animal was allocated in the center of the equipment where it could freely explore the arena for 5 minutes. From this test, the total number of crossings from one quadrant to another (number of line crossings) was measured. A crossing was considered only when all the 4 rat's paws were within the same quadrant. The objects and the box were cleaned with 10% alcohol in order to avoid olfactory bias between each animal; all assays took place in a controlled lighting room (20 lx).

### **2.5.2 Y Maze Test**

The Y Maze test was performed on the 29<sup>th</sup> day after surgery as previously described by Bassani, Turnes [38]. Using this test it was possible to assess short term spatial memory [39, 40]. The Y Maze consists of an apparatus in the shape of the letter Y, made of wood and painted black, with three arms separated by 120° angles, measuring 50 cm in length, 12 cm in width and 27 cm in height. The animals' behavior was filmed for later evaluation with a camera. In the training session of the Y Maze, one of the arms was unreachable by a wooden door located in front of it. The rats were placed in one of the other two arms (initial arm), and were able to explore these maze arms for 5 min. After 1h of the training session it was performed the Y Maze test itself. The blockade was detached, allowing access to all of the arms. The previously blocked arm was named “new arm”. Rats were placed back in the initial arm and were able to freely explore for 3 min. To evaluate the short-term spatial memory, was considered the percentage of time exploring the new arm, regarding the total exploratory time [41]. The maze was cleaned with 10% alcohol between each session and each animal to reduce olfactory bias; all assays took place in a controlled lighting room (20 lx).

### **2.5.3 NORT**

The NORT was performed in order to assess short-term memory [42]. The percentage of exploration of the new object was measured, in relation to the total exploratory time of the two objects [43]. The environment and gears used were standardized by Bassani et al (2017). Test and training (habituation) took place in a square box (100 x 100 x 40 cm), made of wood. A camera was positioned to obtain an aerial view of the arena to record the animals' behavior, for later evaluation. Before NORT was performed, first the animals went through a habituation stage on the 28th day after the surgery, where each animal was placed in the empty box for 5 minutes of free exploration. After that, a new 5 minutes habituation session was held 24h later, on this session 2 identical objects were positioned symmetrically in the apparatus, approximately 10 cm away from the walls. The animals were able to explore the objects for 5 min and then were returned to their housing boxes. Subsequently the second training session a third one was performed, identical to the previous session, respecting an 1h interval. After 1h of the training sessions, the NORT itself was

implemented (29th day after surgery). On NORT one of the objects was kept (familiar object) and the other one was replaced by a different object (new object). Both remained in their original positions. Each animal was placed in the arena for 3 minutes to explore the objects [44]. In order to avoid olfactory bias, before each session and between each animal, the objects and the box were cleaned with 10% alcohol; all assays took place in a controlled lighting room (20 lx).

## **2.6. Neurochemical Analysis**

At the 30th day after surgery, the brains were rapidly extracted on dry ice, and the whole hippocampus and prefrontal cortex were dissected and stored at  $-80^{\circ}\text{C}$ . The tissue homogenate of these biological samples was prepared, according to the manufacturer's instructions (USCN Business Co., Wuhan, Hubei, China) and Enzyme-Linked Immunosorbent Assay (ELISA) was made for  $\text{A}\beta$  and  $\alpha$ -Syn. The tissues were washed in ice-cold phosphate buffered saline (PBS) to remove blood excess and weighed before homogenization. After that, the samples were minced and homogenized in fresh lysis buffer (complete™, EDTA-free Protease Inhibitor Cocktail from Roche). The resulting suspension was sonicated with an ultrasonic cell disrupter to clarification. After that the homogenates were centrifuged for 5 minutes at  $10.000\times g$  and then was collect the supernatant to perform the assay. Subsequently, was performed the quantification of  $\text{A}\beta$  (product No. CEA946Ra) and  $\alpha$ -Syn (product No. SEB222Ra) of these samples by ELISA according to the manufacturer's instructions (USCN Business Co., Wuhan, Hubei, China). The assays were performed as duplicates and the means of the microplate reader were used (measured at 450 nm), its results are expressed as absorbance.

## **2.7. Statistical Analysis**


Data was presented as mean  $\pm$  standard error of mean (SEM). The Kolmogorov-Smirnov test was used to confirm the data was normally distributed. In order to assess the influence of both factors (Galantamine and STZ) on all response variables (behavioral tests and ELISA), as well as factors interactions, difference between groups were assessed performing a two-way Analysis of Variance (ANOVA) with significance level equal to 5% ( $\alpha = 0.05$ ).

Thus, in this paper, when the ANOVA test showed a Fisher's  $p$ -value ( $p$ ) less than the significance level for any source of variation, i.e.  $p < 0.05$ , Tukey's multiple comparisons tests were made to investigate means differences between each group. Here, again, a significant difference was considered when the  $p < 0.05$ .

### 3. Results

#### 3.1. Locomotion Frequency Test

On the OFT it was measured the spontaneous locomotor performance activity by accessing the total number of crossings (Figure 3). The ANOVA test for the OFT only showed statistically significance difference for the STZ effect ( $p = 0.0001$ ). The Tukey test shows statistically significant difference between Sham + Galantamine and STZ + Vehicle groups ( $p = 0.0017$ ). There is also statistically significant difference between Sham + Galantamine and STZ + Galantamine groups ( $p = 0.0050$ ).




**Figure 3** – Effects of prolonged Galantamine (5 mg/kg) in spontaneous locomotor activity of Sham and STZ groups. All data is expressed as mean  $\pm$  SEM.  $*p < 0.05$ , indicating a statistically significant difference between groups. (ANOVA).  $N = 8$  animals per group.



### 3.2. Memory Tests


Through the Y Maze test it was possible to assess the short-term spatial memory, by measuring the percentage of time exploring the new arm (Figure 4).

The ANOVA test for the Y Maze only showed statistically significance difference for the STZ effect ( $p = 0.0041$ ). Multiple comparison Tukey's test showed a statistically significant difference between the Sham + Vehicle and STZ + Vehicle groups ( $p = 0.0218$ ).



**Figure 4** – Prolonged Galantamine administration (5 mg/kg) and its impact in the short-term spatial memory. All data are expressed as mean ± SEM. \* $p < 0.05$ , indicating a statistically significant difference between groups. (ANOVA).  $N = 8$  animals per group.

No effect of treatment was seen in the percentage of exploratory time of the new object (Figure 5).




**Figure 5** – Short-term memory effects of prolonged Galantamine (5 mg/kg) administration in Sham and STZ groups. The data are expressed as mean  $\pm$  SEM.  $N = 8$  animals per group. There was no statistically significant difference between groups (ANOVA).

### 3.3. Quantification of beta amyloid and alpha synuclein by enzyme-linked immunosorbent assay (ELISA)

The accumulation of  $A\beta$  in the prefrontal cortex and in the hippocampus showed an increase in this protein in the STZ-lesioned rats ( $p = 0.0002$  and  $p = 0.0048$  respectively). In cortex both STZ groups presented higher  $A\beta$  absorbance than the Sham groups (Figure 6A). Regarding the Galantamine treatment, there was no difference between STZ + Vehicle and STZ+ Galantamine. In hippocampus there was more accumulation of  $A\beta$  in the STZ groups but the STZ + Galantamine group had less  $A\beta$  concentration, when compared to the STZ + Vehicle group. The only statistically significant difference observed in Figure 6B is between the Sham + Galantamine and STZ + Vehicle groups ( $p = 0.0112$ ).


**Figure 6 - Quantification of  $A\beta$  levels in the prefrontal cortex (A) and hippocampus (B)**



by ELISA. Statistical relationship between animals injured with STZ and Sham. It is also shown the implications of prolonged use of Galantamine (5 mg/kg) in either group. All data is expressed as mean  $\pm$  SEM.  $*p < 0.05$ , indicating a statistically significant difference between groups. (ANOVA).  $N = 8$  animals per group.

When it comes to  $\alpha$ -Syn accumulation, it is observed an accumulation of this protein in the cortex of the STZ groups (Figure 7A). Excluding the statistical difference between Sham + Galantamine and STZ + Vehicle, Galantamine had no impact in the accretion of this protein, regarding cortex.

The phenomenon of more  $\alpha$ -Syn accumulation in the STZ groups is not observed in the hippocampus (Figure 7B), however, when observing the galantamine effect, it is notice less  $\alpha$ -Syn in the STZ + Galantamine group when compared to the STZ + Vehicle group. Moreover, it is observed a statistical difference between the Sham + Galantamine and STZ + Vehicle groups:  $p = 0.0268$ .



**Figure 7** - Quantification of  $\alpha$ -Syn levels in the prefrontal cortex (**A**) and hippocampus (**B**) by ELISA. Statistical relationship between animals injured with STZ and Sham. It is also shown the implications of prolonged use of Galantamine (5 mg/kg) in either group. All data is expressed as mean  $\pm$  SEM. \* $p < 0.05$ , indicating a statistically significant difference between groups. (ANOVA).  $N = 8$  animals per group.

#### 4. Discussion

The present study investigated the STZ-lesion in galantamine treated rats. The results showed that STZ groups had reduced locomotion frequency in addition to a lesser percentage of time exploring the new arm in Y maze test. we found that the STZ increased the accumulation of  $A\beta$  in the prefrontal cortex as well as in the hippocampus. Moreover, the STZ-lesioned rats exhibited an increase in the  $\alpha$ -Syn in the prefrontal cortex. The Galantamine treatment did not present signs of reversal in these effects.

The OFT and Y Maze test showed significant differences in the locomotion frequency and memory of the animals (Figure 3). The performance of the animals in the STZ + Vehicle and STZ + Galantamine groups are similar, indicating that the Galantamine had no effect on the locomotion frequency. Furthermore it is shown that both the Sham groups presented better performance than the STZ groups on the OFT. These results show that the injured animals had a diminished locomotion frequency.

Moreover, as can be seen by the results of the Y Maze test (Figure 4), the mean percentage of time exploring the new arm by the Sham + Vehicle group was significantly higher than the STZ + Vehicle group. One can notice that all STZ groups presented lower memory performance in this test. This result gives support to the AD's animal model, since the short-term spatial memory on STZ + Vehicle is damaged when compared to the Sham + Vehicle group. Furthermore, there was no statistically significant difference between groups treated with the Galantamine.

Regarding the NORT results, it was observed (Figure 5) there were no statistically significant differences between any of the groups. Nonetheless, even without statistical significance, the STZ + Vehicle group had diminished memory performance when compared to all other groups. In addition, the STZ + Galantamine group presented better performance than the STZ + Vehicle group. These results may indicate that the Galantamine could ease the AD symptoms. However, more researches are needed to further investigate this hypothesis.

Since in the Novel Object Recognition behavioral test there were no statistically significant differences between Sham and STZ groups, one could suggest that the STZ animals were not in an advanced AD stage. One possible explanation to this result could be related to the low dose of STZ. What gives substantial support to this hypothesis is that the spatial memory is one of the first types of memory affected by the AD, with some authors even suggesting that it may occur in preclinical stages [45]. In this regard, our results indicate that the STZ group presented impairments on its spatial memory performance, since these animals did not perform well on the Y Maze test (Figure 4).

In this scenario, we suggest that memory and locomotor performances were diminished due to the same trigger mechanism: the IRBS and oxidative stress, since STZ is a drug capable of inducing glucose metabolism impairments through its affinity with GLUT-2 [18]. In this connection, some authors even suggest that AD may be a type 3 diabetes [46, 47].

However, no statistically significant difference regarding memory and locomotion improvement due to the Galantamine treatment was found on the animals impaired by the STZ. Possibly the lack of the Galantamine effect observed in this paper is due to its action mechanism and the temporal window in which this drug is able to produce beneficial outcomes. Galantamine is a selective, competitive and reversible acetylcholinesterase inhibitor.

In addition, it is able to increase the intrinsic action of acetylcholine on nicotinic receptors (modulation), probably through binding to an allosteric site of this receptor [48]. As consequence of the improvement in the cholinergic system, a boost in the cognitive functions of AD patients can be observed [49].

Although acetylcholinesterase inhibitors can temporarily attenuate memory and cognitive impairments in AD patients [49], AD is a progressive disease and currently there is no treatment regarding the ongoing cellular loss [48].

In this line, in 2012, a study in Guinea pigs showed that Galantamine prevents cell death due to apoptosis induced by Soman 28 µg/kg, an anticholinesterase drug [50]. Also, there is indicatives that there is a link regarding the basal forebrain cholinergic system and the AD pathogenesis [51]. Since Galantamine is a reversible acetylcholinesterase inhibitor, being an activator of cholinergic-inflammatory pathway by suppressing levels of TNF and IL-6 [30, 35, 52-54], is supposed that this drug is able to improve the inflammatory cytokines response, consequently interfering in the A $\beta$  formation cascade [55].

In addition, the Galantamine presents disease-modifying capacities, as has been reported in Van Dam and De Deyn [56], and references therein, in this scenario, we suspect that the insufficient performance of the Galantamine in our study is linked with the preventive characteristics of this drug. Perhaps a more prolonged treatment before stereotaxic surgery [57], combined with a different dosage schedule aiming at the preventive and disease-modifying characteristics, would have found a different outcome for the results shown in this paper.

The lesser frequency locomotion found in the rats could be due to the  $\alpha$ -Syn accumulation and not only because of the AD animal model induced by STZ. In addition, another noticeable result is that in most of the tests there were no statistically significant differences in the memory of the rats treated with Galantamine, when compared to those only treated with vehicle. However, one cannot assume promptly that Galantamine has no effect whatsoever in treating AD clinical features, since some differences were observed.

Regarding the A $\beta$  accumulation measured by the ELISA (Figure 6), the STZ + Galantamine group presented a lower absorbance in the prefrontal cortex, when compared to the Sham + Vehicle group (Figure 6A). This result indicates that the STZ reproduces a feature of AD, which was an expected result since this type of AD animal

model was already described by Bassani et al. (2017). Moreover, other differences point to the fact that Galantamine has no effect mitigating the neuroinflammatory cascade, since both STZ groups presented higher  $A\beta$  absorbance in the prefrontal cortex than the Sham groups, as can be seen in Figure 6A.

The results observed for the prefrontal cortex also support the AD STZ animal model (Figure 6B), showing that the STZ impaired rats presented higher  $A\beta$  absorbance than the Sham ones. There was statistically significant difference between the Sham + Galantamine and STZ + Vehicle groups, due to the STZ effect, as shown by the two-way ANOVA result ( $p = 0.0048$ ).

The results of the ELISA with respect to  $\alpha$ -Syn absorbance (Figure 7) indicates that not only the animals had AD biomarkers, but also presented what is usually considered an exclusive PD biomarker [58]. The high absorbance of the  $\alpha$ -Syn in the prefrontal cortex, seen on Figure 7A, may be the cause of the performance impairment seen on the OFT, indicating that the rats could be suffering from parkinsonism.

It is well-known that, in contrast to the AD, PD is characterized by motor impairment caused by accumulation of  $\alpha$ -Syn in dopaminergic neurons, forming structures called Lewy bodies [59]. In addition, in its late stages, PD may cause cognitive deficits and dementia [60-62]. Despite one can find different protein accumulation in the neural tissue of AD and PD patients brains,  $A\beta$  and  $\alpha$ -Syn, respectively, both diseases may have common pathological triggers, like neuroinflammation and oxidative stress [63], as well as both proteins accumulation [16, 64]. Thus, some scientists believe that PD and AD may have a common metabolic physiopathology [21].

Hence, although one can suggest that PD and AD may have common trigger mechanisms, such as oxidative stress [17], the fact that in this paper was found higher  $\alpha$ -Syn absorbance in the STZ lesioned rats, it is yet not enough to confirm that the rats have mimicked the PD thoroughly. We believe that the animals have presented motor impairment due to the  $\alpha$ -Syn accumulation. This statement could be accepted, since some studies demonstrate that up to 60% of patients with AD also present signs of Lewy body disorders [11-14].

Regarding the difference in  $\alpha$ -Syn accumulation in the Hippocampus between the Sham + Vehicle and STZ + Galantamine groups, the two-way ANOVA indicated that the variation between groups was due to the Galantamine effect ( $p = 0.0292$ ). Hence,



the phenomenon described in the previous paragraph could explain the results shown on Figure 7B. The lower absorbance of  $\alpha$ -Syn in the impaired group due to the Galantamine effect indicates that the drug could ease the IRBS and its damages.

Finally, the groups treated with STZ presented both A $\beta$  and  $\alpha$ -Syn accumulation, indicating that the high absorbance of these proteins may have a common source.

Nevertheless, since, to the best of the authors knowledge, this is one of the first studies concerning Galantamine effects on the AD animal model induced by STZ, as well as on the assessment of  $\alpha$ -Syn accumulation, we believe that our results cheer more researches in this direction.

## 5. Concluding remarks

This paper presented the Galantamine effects on a STZ AD animal model. Despite the fact that the animals impaired with STZ did not presented improvements when treated with Galantamine, the overall performance for the STZ + Galantamine group was higher than the STZ + Vehicle one on the NORT. On the other hand, on the ELISA all the STZ groups had higher  $\alpha$ -Syn and A $\beta$  accumulation in their Hippocampus and Prefrontal Cortex. One result (Figure 7B) indicated that Galantamine may have diminished the  $\alpha$ -Syn accumulation of the STZ impaired group.

The locomotion impairment found in the animals is due to the  $\alpha$ -Syn accumulation, and not because of the AD. STZ animals were not in an advanced AD stage, the impairment could be due to the IRBS, which can cause either  $\alpha$ -Syn accumulation and memory impairment (AD). Nevertheless, we believe that more studies on this subject should be carried out to elucidate some challenges yet unresolved.

## Acknowledgements

We would like to thank Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Ensino Superior (CAPES) for the financial support. MABFV is recipient of a National Counsel of Technological and Scientific Development fellowship. The authors report no conflicts of interest.

## References

1. Cipriani, G., et al., *Depression and dementia. A review*. European Geriatric Medicine, 2015. **6**(5): p. 479-486.
2. dos Santos, P., et al., *Alzheimer's disease: a review from the pathophysiology to diagnosis, new perspectives for pharmacological treatment*. Current medicinal chemistry, 2018. **25**(26): p. 3141-3159.
3. Silvestrelli, G., et al., *Treatment of Alzheimer's disease: from pharmacology to a better understanding of disease pathophysiology*. Mechanisms of ageing and development, 2006. **127**(2): p. 148-157.
4. Lowe, V.J., et al., *Widespread brain tau and its association with ageing, Braak stage and Alzheimer's dementia*. Brain, 2018. **141**(1): p. 271-287.
5. Kamat, P.K., et al., *Streptozotocin intracerebroventricular-induced neurotoxicity and brain insulin resistance: A therapeutic intervention for treatment of sporadic Alzheimer's disease (sAD)-like pathology*. Molecular neurobiology, 2016. **53**(7): p. 4548-4562.
6. Pákási, M. and J. Kálmán, *Interactions between the amyloid and cholinergic mechanisms in Alzheimer's disease*. Neurochemistry international, 2008. **53**(5): p. 103-111.
7. Barage, S.H. and K.D. Sonawane, *Amyloid cascade hypothesis: Pathogenesis and therapeutic strategies in Alzheimer's disease*. Neuropeptides, 2015. **52**: p. 1-18.
8. Sereniki, A. and M.A.B.F. Vital, *A doença de Alzheimer: aspectos fisiopatológicos e farmacológicos*. Revista de psiquiatria do Rio Grande do Sul, 2008. **30**(1): p. 0-0.
9. Spillantini, M. and D. Dickson, *Introduction to synucleinopathies*. Neurodegeneration: the molecular pathology of dementia and movement disorders. Basel: ISN Neuropath, 2003: p. 156-8.
10. Chang, C.-W., et al., *Plasma and Serum Alpha-Synuclein as a Biomarker of Diagnosis in Patients With Parkinson's Disease*. Frontiers in Neurology, 2020. **10**: p. 1388.
11. Hamilton, R.L., *Lewy bodies in Alzheimer's disease: a neuropathological review of 145 cases using  $\alpha$ -synuclein immunohistochemistry*. Brain pathology, 2000. **10**(3): p. 378-384.
12. Schneider, J.A., et al., *The neuropathology of probable Alzheimer disease and mild cognitive impairment*. Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society, 2009. **66**(2): p. 200-208.
13. Arai, Y., et al.,  *$\alpha$ -Synuclein-positive structures in cases with sporadic Alzheimer's disease: morphology and its relationship to tau aggregation*. Brain research, 2001. **888**(2): p. 287-296.
14. James, B.D., et al., *Dementia from Alzheimer disease and mixed pathologies in the oldest old*. Jama, 2012. **307**(17): p. 1798-1800.
15. Majbour, N.K., et al., *Increased levels of CSF total but not oligomeric or phosphorylated forms of alpha-synuclein in patients diagnosed with probable Alzheimer's disease*. Scientific reports, 2017. **7**(1): p. 1-8.
16. Mikolaenko, I., et al., *Alpha-synuclein lesions in normal aging, Parkinson disease, and Alzheimer disease: evidence from the Baltimore Longitudinal*

- Study of Aging (BLSA)*. Journal of Neuropathology & Experimental Neurology, 2005. **64**(2): p. 156-162.
17. Angelova, P.R. and A.Y. Abramov, *Alpha-synuclein and beta-amyloid—different targets, same players: calcium, free radicals and mitochondria in the mechanism of neurodegeneration*. Biochemical and biophysical research communications, 2017. **483**(4): p. 1110-1115.
  18. Grieb, P., *Intracerebroventricular streptozotocin injections as a model of Alzheimer's disease: in search of a relevant mechanism*. Molecular neurobiology, 2016. **53**(3): p. 1741-1752.
  19. Steen, E., et al., *Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes?* Journal of Alzheimer's disease, 2005. **7**(1): p. 63-80.
  20. De la Monte, S.M. and J.R. Wands, *Alzheimer's disease is type 3 diabetes—evidence reviewed*. Journal of diabetes science and technology, 2008. **2**(6): p. 1101-1113.
  21. Barilar, J.O., et al., *Shared cerebral metabolic pathology in non-transgenic animal models of Alzheimer's and Parkinson's disease*. Journal of Neural Transmission, 2020: p. 1-20.
  22. Agrawal, R., et al., *A study of brain insulin receptors, AChE activity and oxidative stress in rat model of ICV STZ induced dementia*. Neuropharmacology, 2009. **56**(4): p. 779-787.
  23. Chen, Y., et al., *A non-transgenic mouse model (icv-STZ mouse) of Alzheimer's disease: similarities to and differences from the transgenic model (3xTg-AD mouse)*. Molecular neurobiology, 2013. **47**(2): p. 711-725.
  24. Prickaerts, J., T. Fahrig, and A. Blokland, *Cognitive performance and biochemical markers in septum, hippocampus and striatum of rats after an icv injection of streptozotocin: a correlation analysis*. Behavioural brain research, 1999. **102**(1-2): p. 73-88.
  25. Salkovic-Petrisic, M., et al., *Cerebral amyloid angiopathy in streptozotocin rat model of sporadic Alzheimer's disease: a long-term follow up study*. Journal of neural transmission, 2011. **118**(5): p. 765-772.
  26. Hoyer, S., et al., *Brain Glucose Metabolism Is Controlled by Amplification and Desensitization of the Neuronal Insulin Receptor*. Annals of the New York Academy of Sciences, 1996. **777**(1): p. 374-379.
  27. Salkovic-Petrisic, M., et al., *What have we learned from the streptozotocin-induced animal model of sporadic Alzheimer's disease, about the therapeutic strategies in Alzheimer's research*. Journal of neural transmission, 2013. **120**(1): p. 233-252.
  28. Bloniecki, V., et al., *EFFECTS OF RISPERIDONE AND GALANTAMINE TREATMENT ON ALZHEIMER BIOMARKER LEVELS IN CEREBROSPINAL FLUID*. Alzheimer's & Dementia: The Journal of the Alzheimer's Association, 2017. **13**(7): p. P1047-P1048.
  29. Geerts, H., *Indicators of neuroprotection with galantamine*. Brain research bulletin, 2005. **64**(6): p. 519-524.
  30. Wang, T., et al., *Galantamine reversed early postoperative cognitive deficit via alleviating inflammation and enhancing synaptic transmission in mouse hippocampus*. European journal of pharmacology, 2019. **846**: p. 63-72.
  31. El-Abhar, H., et al., *Galantamine: anti-Diabetic Effect Mediated By Anti-Oxidant/Inflammatory/Apoptotic Effects And Improvement of Insulin And Wnt/ $\beta$ -Catenin Signaling Pathways*. Clinical Therapeutics, 2015. **37**(8): p. e31.

32. Ali, M.A., et al., *Antidiabetic effect of galantamine: novel effect for a known centrally acting drug*. PloS one, 2015. **10**(8).
33. Ali, M.A.I., *Potential Antidiabetic Effect of Galantamine, a Known Centrally Acting Drug with or without Vildagliptin in Type 2 Diabetic Rats*. CU Theses, 2015.
34. Furukawa, S., L. Yang, and H. Sameshima, *Galantamine, an acetylcholinesterase inhibitor, reduces brain damage induced by hypoxia-ischemia in newborn rats*. International Journal of Developmental Neuroscience, 2014. **37**: p. 52-57.
35. Satapathy, S.K., et al., *Galantamine alleviates inflammation and other obesity-associated complications in high-fat diet-fed mice*. Molecular medicine, 2011. **17**(7-8): p. 599-606.
36. Geerts, H., et al., *Brain levels and acetylcholinesterase inhibition with galantamine and donepezil in rats, mice, and rabbits*. Brain research, 2005. **1033**(2): p. 186-193.
37. Paxinos, G. and C. Watson, *The rat brain in stereotaxic coordinates: hard cover edition*. 2006: Elsevier.
38. Bassani, T.B., et al., *Effects of curcumin on short-term spatial and recognition memory, adult neurogenesis and neuroinflammation in a streptozotocin-induced rat model of dementia of Alzheimer's type*. Behavioural brain research, 2017. **335**: p. 41-54.
39. Conrad, C.D., et al., *Chronic stress impairs rat spatial memory on the Y maze, and this effect is blocked by tianeptine treatment*. Behavioral neuroscience, 1996. **110**(6): p. 1321.
40. Dean, O., et al., *Glutathione depletion in the brain disrupts short-term spatial memory in the Y-maze in rats and mice*. Behavioural brain research, 2009. **198**(1): p. 258-262.
41. Huang, H., et al., *Isolation housing exacerbates Alzheimer's disease-like pathophysiology in aged APP/PS1 mice*. International Journal of Neuropsychopharmacology, 2015. **18**(7): p. pyu116.
42. Antunes, M. and G. Biala, *The novel object recognition memory: neurobiology, test procedure, and its modifications*. Cognitive processing, 2012. **13**(2): p. 93-110.
43. Walf, A.A., M.E. Rhodes, and C.A. Frye, *Ovarian steroids enhance object recognition in naturally cycling and ovariectomized, hormone-primed rats*. Neurobiology of learning and memory, 2006. **86**(1): p. 35-46.
44. De Bruin, N., et al., *Two novel 5-HT<sub>6</sub> receptor antagonists ameliorate scopolamine-induced memory deficits in the object recognition and object location tasks in Wistar rats*. Neurobiology of learning and memory, 2011. **96**(2): p. 392-402.
45. Johnson, D.K., et al., *Longitudinal study of the transition from healthy aging to Alzheimer disease*. Archives of neurology, 2009. **66**(10): p. 1254-1259.
46. Suzanne, M., *Type 3 diabetes is sporadic Alzheimer's disease: mini-review*. European Neuropsychopharmacology, 2014. **24**(12): p. 1954-1960.
47. Hoyer, S., *Glucose metabolism and insulin receptor signal transduction in Alzheimer disease*. European journal of pharmacology, 2004. **490**(1-3): p. 115-125.
48. Arias, E., et al., *Galantamine prevents apoptosis induced by  $\beta$ -amyloid and thapsigargin: involvement of nicotinic acetylcholine receptors*. Neuropharmacology, 2004. **46**(1): p. 103-114.



49. Bonner, L.T. and E.R. Peskind, *Pharmacologic treatments of dementia*. The Medical Clinics of North America, 2002. **86**(3): p. 657-674.
50. Kulkarni, G.S., *Galantamine prevents soman-induced apoptotic cell death in the guinea pig brain*. 2012.
51. Auld, D.S., et al., *Alzheimer's disease and the basal forebrain cholinergic system: relations to  $\beta$ -amyloid peptides, cognition, and treatment strategies*. Progress in neurobiology, 2002. **68**(3): p. 209-245.
52. Hanes, W.M., et al., *Galantamine attenuates type 1 diabetes and inhibits anti-insulin antibodies in nonobese diabetic mice*. Molecular Medicine, 2015. **21**(1): p. 702-708.
53. Martín, A., M. Domercq, and C. Matute, *Inflammation in stroke: the role of cholinergic, purinergic and glutamatergic signaling*. Therapeutic advances in neurological disorders, 2018. **11**: p. 1756286418774267.
54. Konrath, E.L., et al., *Alkaloids as a source of potential anticholinesterase inhibitors for the treatment of Alzheimer's disease*. Journal of Pharmacy and Pharmacology, 2013. **65**(12): p. 1701-1725.
55. Liu, Y., et al., *Galantamine improves cognition, hippocampal inflammation, and synaptic plasticity impairments induced by lipopolysaccharide in mice*. Journal of neuroinflammation, 2018. **15**(1): p. 112.
56. Van Dam, D. and P.P. De Deyn, *Cognitive evaluation of disease-modifying efficacy of galantamine and memantine in the APP23 model*. European neuropsychopharmacology, 2006. **16**(1): p. 59-69.
57. Liu, X., et al., *Protective effects of galantamine against A $\beta$ -induced PC12 cell apoptosis by preventing mitochondrial dysfunction and endoplasmic reticulum stress*. Neurochemistry international, 2010. **57**(5): p. 588-599.
58. Atik, A., T. Stewart, and J. Zhang, *Alpha-synuclein as a biomarker for Parkinson's disease*. Brain pathology, 2016. **26**(3): p. 410-418.
59. Davie, C.A., *A review of Parkinson's disease*. British medical bulletin, 2008. **86**(1): p. 109-127.
60. Gratwicke, J., M. Jahanshahi, and T. Foltynie, *Parkinson's disease dementia: a neural networks perspective*. Brain, 2015. **138**(6): p. 1454-1476.
61. Morris, J.C., et al., *Clinical and pathological aspects of parkinsonism in Alzheimer's disease: a role for extranigral factors?* Archives of neurology, 1989. **46**(6): p. 651-657.
62. Appel, S.H., *A unifying hypothesis for the cause of amyotrophic lateral sclerosis, parkinsonism, and Alzheimer disease*. Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society, 1981. **10**(6): p. 499-505.
63. Han, Z., et al., *Parkinson's disease and Alzheimer's disease: a Mendelian randomization study*. BMC medical genetics, 2018. **19**(1): p. 215.
64. Jellinger, K.A.,  *$\alpha$ -Synuclein pathology in Parkinson's and Alzheimer's disease brain: incidence and topographic distribution—a pilot study*. Acta neuropathologica, 2003. **106**(3): p. 191-202.

#### 4 CONSIDERAÇÕES FINAIS

Correspondendo de 60% a 80% dos quadros demenciais (Götz *et al.*, 2018), a DA ainda representa um grande desafio tanto do ponto de vista do entendimento da doença, quanto do seu tratamento. Esses dois fatores devem servir como estímulo a mais pesquisas nessa linha, pois, atualmente, é uma das patologias mais temidas, que carece de terapêutica modificadora da doença, bem como curativa. A presente tese demonstrou a possibilidade da DA possuir fatores fisiopatológicos em comum com a DP, o que demonstra que nem ao menos a completa história natural da doença dominamos. Diante desse quadro é fácil entender o porquê de ainda não existirem tratamentos eficazes, que curem e ou modifiquem os sombrios desfechos clínicos encontrados na DA.

O modelo animal da DA pela STZ se mostrou eficaz para avaliar os efeitos da Galantamina, produzindo características clínicas e bioquímicas condizentes com a doença. Ainda não existindo consenso sobre os gatilhos iniciadores da formação da  $A\beta$ , as crescentes evidências da presença de  $\alpha$ -Syn despertam ainda mais enigmas sobre a DA, ao mesmo tempo que proporcionam potenciais alvos terapêuticos para a intervenção nessa patologia.

Embora nesse trabalho a Galantamina tenha demonstrado efeitos menores nos sinais correspondentes à DA nos animais. Esse fármaco apresenta relativamente bons resultados no manejo clínico de pacientes com DA moderada a grave. Enquanto nenhuma outra terapia com eficácia comparável ou melhor estiver disponível, a Galantamina ainda permanecerá como uma droga de primeira escolha para o tratamento da DA (Prvulovic *et al.*, 2010).

## 5 CONCLUSÃO

O artigo dessa tese demonstrou efeitos da Galantamina em um modelo animal de DA induzido por STZ. Apesar de os animais lesados pela STZ não apresentarem melhora quando tratados com Galantamina, o desempenho do grupo STZ + Galantamina foi superior ao do grupo STZ + Veículo no Teste de Reconhecimento de Novo Objeto.

Todos os grupos STZ apresentaram maior acúmulo de  $\alpha$ -Syn e A $\beta$  em seu Hipocampo e córtex pré-frontal. No Hipocampo, a Galantamina pode ter diminuído o acúmulo de  $\alpha$ -Syn nos ratos lesados pela STZ.

Acreditamos que o comprometimento da locomoção encontrado nos animais é decorrente do acúmulo de  $\alpha$ -Syn e não apenas de A $\beta$ , indicando que a DA pode ser mais complexa do que o modelo clássico que apenas considera o acúmulo de A $\beta$  como fator patognomônico.

Sugerimos que os animais lesados pela STZ não estavam em estágio avançado da DA, uma vez que não houveram diferenças estatisticamente significantes entre os grupos no Teste de Reconhecimento de Novo Objeto.

Por fim, propomos que a DA e a DP podem ter origem comum, o estado cerebral de resistência à insulina, pelo fato de os grupos tratados com STZ terem apresentado acúmulo tanto de A $\beta$  quanto de  $\alpha$ -Syn.



## REFERÊNCIAS

- ALI, M. A. I. Potential Antidiabetic Effect of Galantamine, a Known Centrally Acting Drug with or without Vildagliptin in Type 2 Diabetic Rats. **CU Theses**, 2015.
- ANGELOVA, P. R.; ABRAMOV, A. Y. Alpha-synuclein and beta-amyloid—different targets, same players: calcium, free radicals and mitochondria in the mechanism of neurodegeneration. **Biochemical and biophysical research communications**, v. 483, n. 4, p. 1110-1115, 2017. ISSN 0006-291X.
- ARAI, Y. et al.  $\alpha$ -Synuclein-positive structures in cases with sporadic Alzheimer's disease: morphology and its relationship to tau aggregation. **Brain research**, v. 888, n. 2, p. 287-296, 2001. ISSN 0006-8993.
- BARAGE, S. H.; SONAWANE, K. D. Amyloid cascade hypothesis: Pathogenesis and therapeutic strategies in Alzheimer's disease. **Neuropeptides**, v. 52, p. 1-18, 2015. ISSN 0143-4179.
- BLONIECKI, V. et al. EFFECTS OF RISPERIDONE AND GALANTAMINE TREATMENT ON ALZHEIMER BIOMARKER LEVELS IN CEREBROSPINAL FLUID. **Alzheimer's & Dementia: The Journal of the Alzheimer's Association**, v. 13, n. 7, p. P1047-P1048, 2017. ISSN 1552-5260.
- CHANG, C.-W. et al. Plasma and Serum Alpha-Synuclein as a Biomarker of Diagnosis in Patients With Parkinson's Disease. **Frontiers in Neurology**, v. 10, p. 1388, 2020. ISSN 1664-2295.
- CIPRIANI, G. et al. Depression and dementia. A review. **European Geriatric Medicine**, v. 6, n. 5, p. 479-486, 2015. ISSN 1878-7649.
- DE LA MONTE, S. M.; WANDS, J. R. Alzheimer's disease is type 3 diabetes—evidence reviewed. **Journal of diabetes science and technology**, v. 2, n. 6, p. 1101-1113, 2008. ISSN 1932-2968.
- DOS SANTOS, P. et al. Alzheimer's disease: a review from the pathophysiology to diagnosis, new perspectives for pharmacological treatment. **Current medicinal chemistry**, v. 25, n. 26, p. 3141-3159, 2018. ISSN 0929-8673.
- EL-ABHAR, H. et al. Galantamine: anti-Diabetic Effect Mediated By Anti-Oxidant/Inflammatory/Apoptotic Effects And Improvement of Insulin And Wnt/ $\beta$ -Catenin Signaling Pathways. **Clinical Therapeutics**, v. 37, n. 8, p. e31, 2015. ISSN 0149-2918.
- FURUKAWA, S.; YANG, L.; SAMESHIMA, H. Galantamine, an acetylcholinesterase inhibitor, reduces brain damage induced by hypoxia-ischemia in newborn rats. **International Journal of Developmental Neuroscience**, v. 37, p. 52-57, 2014. ISSN 0736-5748.

- GEERTS, H. Indicators of neuroprotection with galantamine. **Brain research bulletin**, v. 64, n. 6, p. 519-524, 2005. ISSN 0361-9230.
- GEERTS, H. et al. Brain levels and acetylcholinesterase inhibition with galantamine and donepezil in rats, mice, and rabbits. **Brain research**, v. 1033, n. 2, p. 186-193, 2005. ISSN 0006-8993.
- GÖTZ, J.; BODEA, L.-G.; GOEDERT, M. Rodent models for Alzheimer disease. **Nature Reviews Neuroscience**, p. 1, 2018. ISSN 1471-0048.
- GRIEB, P. Intracerebroventricular streptozotocin injections as a model of Alzheimer's disease: in search of a relevant mechanism. **Molecular neurobiology**, v. 53, n. 3, p. 1741-1752, 2016. ISSN 0893-7648.
- HAMILTON, R. L. Lewy bodies in Alzheimer's disease: a neuropathological review of 145 cases using  $\alpha$ -synuclein immunohistochemistry. **Brain pathology**, v. 10, n. 3, p. 378-384, 2000. ISSN 1015-6305.
- JAMES, B. D. et al. Dementia from Alzheimer disease and mixed pathologies in the oldest old. **Jama**, v. 307, n. 17, p. 1798-1800, 2012. ISSN 0098-7484.
- KAMAT, P. K. et al. Streptozotocin intracerebroventricular-induced neurotoxicity and brain insulin resistance: A therapeutic intervention for treatment of sporadic Alzheimer's disease (sAD)-like pathology. **Molecular neurobiology**, v. 53, n. 7, p. 4548-4562, 2016. ISSN 0893-7648.
- LOWE, V. J. et al. Widespread brain tau and its association with ageing, Braak stage and Alzheimer's dementia. **Brain**, v. 141, n. 1, p. 271-287, 2018. ISSN 0006-8950.
- MAJBOUR, N. K. et al. Increased levels of CSF total but not oligomeric or phosphorylated forms of alpha-synuclein in patients diagnosed with probable Alzheimer's disease. **Scientific reports**, v. 7, n. 1, p. 1-8, 2017. ISSN 2045-2322.
- MIKOLAENKO, I. et al. Alpha-synuclein lesions in normal aging, Parkinson disease, and Alzheimer disease: evidence from the Baltimore Longitudinal Study of Aging (BLSA). **Journal of Neuropathology & Experimental Neurology**, v. 64, n. 2, p. 156-162, 2005. ISSN 1554-6578.
- PÁKÁSKI, M.; KÁLMÁN, J. Interactions between the amyloid and cholinergic mechanisms in Alzheimer's disease. **Neurochemistry international**, v. 53, n. 5, p. 103-111, 2008. ISSN 0197-0186.
- PRVULOVIC, D.; HAMPEL, H.; PANTEL, J. Galantamine for Alzheimer's disease. **Expert opinion on drug metabolism & toxicology**, v. 6, n. 3, p. 345-354, 2010. ISSN 1742-5255.

- SATAPATHY, S. K. et al. Galantamine alleviates inflammation and other obesity-associated complications in high-fat diet-fed mice. **Molecular medicine**, v. 17, n. 7-8, p. 599-606, 2011. ISSN 1076-1551.
- SCHNEIDER, J. A. et al. The neuropathology of probable Alzheimer disease and mild cognitive impairment. **Annals of Neurology: Official Journal of the American Neurological Association and the Child Neurology Society**, v. 66, n. 2, p. 200-208, 2009. ISSN 0364-5134.
- SERENIKI, A.; VITAL, M. A. B. F. A doença de Alzheimer: aspectos fisiopatológicos e farmacológicos. **Revista de psiquiatria do Rio Grande do Sul**, v. 30, n. 1, p. 0-0, 2008. ISSN 0101-8108.
- SILVESTRELLI, G. et al. Treatment of Alzheimer's disease: from pharmacology to a better understanding of disease pathophysiology. **Mechanisms of ageing and development**, v. 127, n. 2, p. 148-157, 2006. ISSN 0047-6374.
- SPELLANTINI, M.; DICKSON, D. Introduction to synucleinopathies. **Neurodegeneration: the molecular pathology of dementia and movement disorders**. Basel: ISN Neuropath, p. 156-8, 2003.
- STEEN, E. et al. Impaired insulin and insulin-like growth factor expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes? **Journal of Alzheimer's disease**, v. 7, n. 1, p. 63-80, 2005. ISSN 1387-2877.
- WANG, T. et al. Galantamine reversed early postoperative cognitive deficit via alleviating inflammation and enhancing synaptic transmission in mouse hippocampus. **European journal of pharmacology**, v. 846, p. 63-72, 2019. ISSN 0014-2999.